

RESEARCH

Open Access



Poor sleep quality and associated factors among HIV-positive pregnant women in Northwest, Ethiopia: a facility-based, cross-sectional study

Getasew Legas^{1*}, Getnet Mihretie Beyene¹, Sintayehu Asnakew¹, Amsalu Belete¹ and Tigabu Desie²

Abstract

Background: Poor sleep quality during pregnancy might have an impact on adverse birth outcomes like premature rupture of membrane, preterm birth, lifelong neurocognitive impairment, low birth weight, and increased the risk of neonatal morbidity and mortality. In Ethiopia, the magnitude of poor sleep quality among this group of people is extremely limited. So, this study aims to determine the magnitude of poor sleep quality and its associated factors among HIV-positive pregnant women attending public hospitals in Northwest Ethiopia.

Methods: An institution-based cross-sectional study was done using a simple random sampling technique to recruit 411 HIV-positive pregnant women from January to March; 2021. Sleep quality over the last 1 month was measured using the Pittsburgh Sleep Quality Index (PSQI). General anxiety disorder (GAD-7), Sleep Hygiene Index (SHI), and List of Threatening of Experiences (LTE) instruments were used to identify factors associated with poor sleep quality. Bivariate and multivariable logistic regression with odds ratio and 95% CI were employed to identify determinant factors of poor sleep quality. Statistical significance association was declared at P -value < 0.05 .

Results: A total of 411 out of 423 HIV-positive pregnant women were interviewed, with a response rate of 97.1%. The overall magnitude of poor sleep quality among HIV-positive pregnant was found to be 39.4% with a 95% of confidence interval (CI) (34.3, 44.3). Stressful life events, [AOR = 3.10, 95% CI (1.60, 6.01)], having comorbid general anxiety symptoms [AOR = 2.46, 95% CI (1.58, 3.81)], unplanned pregnancy [AOR = 2.18, 95% CI (1.20, 3.96)], and poor sleep hygiene practice [AOR = 2.23, 95% CI (1.21, 4.10)] were significantly associated with poor quality of sleep.

Conclusion: The overall magnitude of poor sleep quality among HIV-positive pregnant women was high. Stressful life events, poor sleep hygiene, unplanned pregnancy, and comorbid general anxiety symptoms were the determinant factors of poor sleep quality that should be taken high consideration for early detection and appropriate intervention for poor sleep quality in HIV-positive pregnant women.

Keywords: Poor sleep quality, HIV-positive, Northwest, Ethiopia

Background

Sleep quality is defined as an individual's satisfaction of sleep with different aspects of the sleep experience including sleep efficiency, sleep latency, sleep duration, and wake after sleep onset [1]. Sleep is a universal behavior, occupying one-third of human life and a

*Correspondence: getasewlegas@gmail.com

¹ Department of Psychiatry, College of Medicine and Health Sciences, Debre Tabor University, Debre Tabor, Ethiopia

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

significant contributor to energy restoration for proper functioning. Prolonged sleep deprivation might be lead to severe physical illness, the development of mental illnesses, substance use disorders, and cognitive impairment [2]. Globally, over 36.9 million people were living with human immunodeficiency virus/Acquired immunodeficiency syndrome (HIV/AIDS) and sub-Saharan countries are the most severely affected region in Africa estimated at one in every 25 people living with HIV/AIDS [3, 4].

According to the reports of different studies, 50 to 70 million people were suffering from sleep disturbance in the United States [5]. Poor quality of sleep is one of the most common sleep disturbances in HIV/AIDS patients, it accounts for 40–70% of individuals suffering from sleep disturbance compared with the general population [6–8]. Poor sleep quality is a common problem in HIV-positive individuals due to direct effect of the virus and opportunistic infection of the central nervous system [9–11].

Sleep disturbance is also common in pregnant women; approximately, up to 78% of women have complained of different forms of sleep disturbance during their pregnancy [12]. It is a major public health problem both in developed and developing countries, and poor sleep quality during pregnancy is a significant contributor to adverse birth outcomes, growth restriction of the fetus [13], including premature rupture of membrane [14], low birth weight, pre-eclampsia [15], prolonged labor [16], stillbirth [17], abruption of placenta [18], gestational diabetes mellitus [19], gestational hypertension [20], pre-term birth [21–24], postpartum depression [25–27], and decreased progesterone level [28].

Studies are carried out to assess the prevalence of poor sleep quality among pregnant women varied across the world. A cross-sectional survey data in China showed that 15.2% of pregnant women had complained of problems related to sleep quality [29]. The other systematic and meta-analysis data in this country also showed the pooled prevalence of poor sleep quality among pregnant women was 54.2% [30]. A study conducted in Bangladesh reported that the magnitude of sleep disturbance was 38.8% [31]. In the USA, 53–71% of pregnant women suffered from poor sleep quality [32], in Vietnam 41.2% [33], in Finland 15% [34], in Malaysia 69.4% [35], in Nepal 28.2% [36], in Taiwan 60% [37], and in Peru 17% [38].

Various factors were identified as a factor of poor sleep quality in the previous studies conducted across the globe, history of abortion, comorbid depression and anxiety [39–42], gestational age (second and third trimester) [43], history of stillbirth, multigravida [44, 45], individual-level of stress [44], stressful life events [41, 46, 47], living in rural areas [48], poor sleep hygiene [42], absence of postnatal care, current use tobacco and khat

[49] were the determinant factors of poor sleep quality during pregnancy.

In Ethiopia, the magnitudes of poor sleep quality in pregnant women range from 30.8 to 68.4% [43–45]. Where poor sleep quality is high in low-and-middle-income countries including Ethiopia. But limited studies were conducted to assess the magnitude of poor sleep quality among HIV-positive pregnant women in Ethiopia. Therefore, assessing the magnitude and associated factors of poor sleep quality among these groups of women is important for early identification, and intervention; to reduce the effect of poor sleep quality on pregnancy and adverse birth outcomes.

Methods

Study design and period

An institution-based cross-sectional study was conducted to determine the magnitude and associated factors of poor sleep quality among HIV-positive pregnant women in Northwest Ethiopia from January to March 2021.

Study area

The study was conducted in the Amhara region selected referral hospitals. The capital city of the Amhara region is Bahirdar which is located northwest part of Ethiopia. The city is the seat of the regional government and is 564 km far from the capital city of Ethiopia (Addis Ababa). The Amhara region is divided into 13 zones named West Gojjam, Awi Zone, North Shewa, Central Gondar, South Gondar, North Gondar, East Gondar, East Gojjam, Oromia zone, Waghimra, South Wollo, North Wollo, and Bahirdar (special zone). According to the recent population census report; the total population size of the Amhara region is estimated at 17,221,976. Of these, 12.27% of the population is residing in the urban area.

Source population

All HIV-positive pregnant women whose age was 18 years and above come for PMTCT (prevention of mothers-to-child-transmission) follow-up in selected Amhara region referral hospitals, Northwest, Ethiopia.

Study population

All HIV-positive pregnant women whose age was 18 years and above come for PMTCT follow-up in selected Amhara region referral hospitals during the study period.

Sampling procedure and sample size determination

A single population proportion formula was used to determine the sample size by taking the prevalence of poor sleep quality among HIV-positive pregnant women was 50% (due to the reason of unknown prevalence) with

a 5% margin of error, and 95% confidence with a 10% non-response rate; the final sample size was 423.

The study subjects were recruited with a simple random sampling technique. The selected referral hospitals in the Amhara region were Woldia, Debre Tabor, Gondar, and Felege Hiwot. In each of the selected hospitals, the number of study participants was obtained from health care providers working in the prevention of mothers-to-child-transmission (PMTCT) clinic. The list of study subjects from each hospital was used as a sampling frame. Then, each study participant was selected by the lottery method.

Inclusion and exclusion criteria

All HIV-positive pregnant women whose age was 18 years and above come for PMTCT follow-up in selected Amhara region referral hospitals were included in the study during the data collection period. HIV-positive pregnant women who were unable to communicate due to severe illness were excluded from the study.

Study variables

Dependent variable

Poor sleep quality (1 = yes, No = 0).

Independent variables

Socio-demographic variables Age, ethnicity, educational status, marital status, and residence.

Clinical factors family history of mental illness, history of stillbirth, sleep hygiene practice, and comorbid general anxiety symptoms.

Psychosocial factors stressful life events.

Operational definition

Sleep quality measured using the Pittsburgh sleep quality index (PSQI) at a cut-off point greater than 5 considered poor sleep quality [50].

Anxiety assessed using GAD-7, a score of ≥ 10 having general anxiety symptoms [51].

Poor sleep hygiene measured using a sleep hygiene index (SHI), a score above the mean on the scale was categorized as poor sleep hygiene practice [52].

Stressful life events The presence of stressful life events in the last 6 months was assessed by using a 12-item List of Threatening Experiences (LTE). A study participant

having one or more stressful life events is considered as having stressful life events [53].

Family history of mental illness was assessed by asking a semi-structured question “Did you have a family history of mental illness?” with a positive response of “Yes” categorized by having a family history of mental illness.

History of stillbirth was measured by asking “Have you had a history of stillbirth?” with a positive response of “yes” considering having a previous history of stillbirth.

Data collection procedures and instruments

A semi-structured interviewer-administered questionnaire was used that contains socio-demographic factors, psychosocial factors (stressful life events, history of stillbirth, and unplanned pregnancy) clinical factors (trimester, family history of mental illness, and comorbid general anxiety symptoms).

Sleep quality was measured using a 19-item of Pittsburgh Sleep Quality Index (PSQI). The tool is categorized into seven components containing subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, daytime dysfunction, and use of sleep medications. Each item of PSQI components has been rated on a three-point scale (0–3), ranging from 0 to 21. A total score of 5 or more according to the PSQI questionnaire was considered as having poor sleep quality over the last month. The tool was validated in Ethiopia with sensitivity and specificity results of 89.6% and 86.5% respectively [50].

General anxiety symptoms in the last 2 weeks were measured using the General Anxiety Disorder questionnaire (GAD-7). The instrument has seven items and each item of questions is rated on a three-point scale, 0 (not at all), 1 (Several days), 2 (More than half the days), and 3 (Nearly every day) with a total score of 0–21. A score of five indicates mild anxiety symptoms, ten indicates moderate anxiety symptoms and fifteen indicates severe symptoms of anxiety [51].

The presence of stressful life events in the last 6 months was assessed by using a 12-item List of Threatening Experiences (LTE). Each item of the instrument has been rated on a two-point scale (yes/no), “1” and “0” indicates the presence and absence of individual level of stress over the last 6 months respectively [53].

Sleep hygiene was measured by a 13-item of sleep hygiene index (SHI). The tool measured the practice of sleep hygiene behavior and each item SHI instrument was rated on a five-point Likert scale characterized by 0(never), 1(rarely), 2(sometimes), 3(frequent), and

4(always) with a total score of 0 to 52. A scoring of above the mean indicates poor sleep hygiene [52].

To examine a family history of mental illness, study subjects were asked: “Did you have a family history of mental illness?” with a response of “yes” having a family history of mental illness. To assess the previous history of stillbirth, study subjects were asked: “Have you had a history of stillbirth?” with a response of “yes” considered a previous history of stillbirth.

Data quality control issues

The data was collected by four trained BSc holder nurses and supervised by trained psychiatry professionals after 2 days of training. The data quality was ensured by giving training on the data collection instrument, sampling procedure, and regular supervision. The questionnaire was also translated from English to the Amharic language (local working language).

Data processing and analysis

The collected data were entered into Epi-data 3.1 and then exported to SPSS - version 21 for analysis. The bivariate logistic analysis was applied to select determinant factors. All independent variables with a p -value < 0.25 in the bivariate analysis were a candidate to enter the multivariable logistic regression model to control the confounding effect. Multivariable logistic regression was used to identify factors associated with poor sleep quality and p -value < 0.05 with 95% of CI considered statistically significant.

Results

A total of 411 out of 423 HIV-positive pregnant women were interviewed, with a response rate of 97.1%. The majority of respondents 380 (92.5%) were Amhara. Half of the study subjects, 206 (50.1%) were in the age group of 18–30 years; regarding religion, almost three-fourth of study participants, 284 (69.1%) were orthodox Christian followers; 363 (86.6%) were currently married, Over half study the respondents, 236 (57.4%) of respondents were urban areas residents, and 255(62%) of respondents were married (Table 1).

Clinical and psychosocial factors of HIV-positive pregnant women

Of the total study subjects, 359 (87.3%) respondents hadn't had stressful life events, and three hundred sixty-nine, (89.8%) hadn't a history of stillbirth. Regarding comorbid mental illness, 268 (65.2%) of respondents had no comorbid anxiety symptoms, 379 (92.2%) of study subjects hadn't a family history of mental illness, and over three-fourth of study participants, 333(81.0%) had good sleep hygiene practice (Table 2).

Table 1 Socio-demographic characteristics of respondents in Northwest Ethiopia, 2021

Variable	Category	Frequency	Percentage
Age	18–30	206	50.1%
	31–45	205	49.9%
Ethnicity	Amhara	380	92.5%
	other	31	7.5%
Educational status	Unable to read and write	95	23.1%
	1–8 grade	127	30.9%
	9–12 grade	63	15.3%
	Diploma & above	126	30.7%
Religion	Orthodox	284	69.1%
	Others	127	30.9%
Marital status	Currently married	355	86.4%
	Currently not married	56	13.6%
Residence	Rural	175	42.6%
	Urban	236	57.4%

Table 2 Distribution of clinical and psychosocial factors of respondents in Northwest, Ethiopia, 2021

Variable	Category	Frequency	Percentage
Unplanned pregnancy	yes	63	15.3
	no	348	84.7
Co-morbid anxiety	yes	143	34.8
	no	268	65.2
History of still birth	Yes	42	10.2
	No	369	89.8
Stressful life events	Yes	52	12.7
	No	359	87.3
Sleep hygiene	good	333	81
	Poor	78	19
Trimester	I	94	22.9
	II	143	34.8
	III	174	42.3
Family history of mental illness	Yes	32	7.8
	No	379	92.2

The magnitudes of poor sleep quality

The overall magnitude of poor sleep quality among HIV-positive pregnant was found to be 39.4% with a 95% of confidence interval (CI) (34.3, 44.3).

Factors associated with poor sleep quality in HIV-positive pregnant women

In this study, poor sleep hygiene, history of stillbirth, unplanned pregnancies, stressful life events, and comorbid anxiety were significantly associated with poor sleep quality on bivariate analysis. In the final adjusted

multivariable analysis, comorbid general anxiety symptoms, poor sleep hygiene, stressful life events, and unplanned pregnancies were found to be significantly associated with poor sleep quality.

The odds of developing poor sleep quality among HIV-positive pregnant women who had unplanned pregnancies were 2-times higher compared with individuals who had planned pregnancies [AOR=2.18, 95% CI (1.20, 3.96)]. Study subjects who had Stressful life events were 3 times more likely to have poor sleep quality than women who hadn't a stressful life event [AOR=3.10, 95% CI (1.60, 6.01)]. The odds of developing poor sleep quality among pregnant women who had poor sleep hygiene was 2-fold times higher than women who had good sleep hygiene [AOR=2.23, 95% CI (1.21, 4.10)]. The likelihood of developing poor sleep quality was 2.46 times higher among study participants who had comorbid general anxiety symptoms compared with those who hadn't comorbid general anxiety symptoms [AOR=2.46, 95% CI (1.58, 3.81)] (Table 3).

Discussion

As per our knowledge, this is the first study to assess poor sleep quality among HIV-positive pregnant women in Africa. In this study, the overall magnitude of poor sleep quality among HIV-positive pregnant was found to be 39.4% with a 95% of confidence interval (CI) (34.3, 44.3). The magnitude of poor sleep quality in the current study was consistent with a study conducted in Bangladesh at 38.8% [31], and Vietnam 41.2% [33]. However, the magnitude of poor sleep quality in the current study is lower than studies reported in Malaysia 69.4% [35], in Taiwan 60% [37], in the USA 53 to 71% [32], in Northern Ethiopia 68.4% [43], and a systematic and meta-analysis data in China which was 54.2% [30].

Conversely, the current study was higher than the estimated magnitude of poor sleeping quality in China 15.2% [29] and 34.14% [14], in Finland 15% [34], and in Southwest Ethiopia 30.8% [44]. The possible reason for the discrepancy might be variations in instrumental used to measure the quality of sleep in HIV-positive pregnant women and variations in the number of sample sizes, cultural, and economic differences might have a significant contribution to this discrepancy.

In the present study, the odds of developing poor sleep quality were 3 times higher among HIV-positive pregnant women who had stressful life events compared with individuals who hadn't stressful life events. This finding was in line with studies done in China [46, 47]. This is explained by the fact that stressful life events had their own impact on the maladaptive hypothalamic-pituitary-adrenal (HPA) system [54]. Negative life events might lead to greater activation of the locus coeruleus norepinephrine system and increased HPA axis, which may also increase excitement and precipitate poor quality of sleep [55].

HIV-positive pregnant women who had unplanned pregnancies were 2-times more likely to have poor sleep quality compared with pregnant women who had planned pregnancies. But, the finding of this study has no similar literature that showed unplanned pregnancies is significantly associated with poor sleep quality. The possible reason might be that HIV-positive pregnant women who had unplanned pregnancy might have psychological and emotional distress. Due to this reason, women who had unplanned pregnancies might have excessive worries. As a result of this, unplanned pregnancies might be a risk factor for poor sleep quality.

In the present study, participants with poor sleep hygiene practice were more likely to have a poor quality

Table 3 Bivariate and Multivariable analyses of poor sleep quality among HIV positive pregnant women in Northwest Ethiopia, 2021

Variables	Category	Sleep quality		COR (95% CI)	AOR (95% CI)
		Good	Poor		
Unplanned pregnancy	No	223	125	1	1
	Yes	26	37	2.53(1.46, 4.38)	2.18 ((1.20, 3.96))*
Comorbid anxiety	No	182	86	1	1
	Yes	67	76	2.40(1.58, 3.64)	2.46 (1.58, 3.81)*
Sleep hygiene	Good	216	117	1	1
	Poor	33	45	2.51(1.53, 4.16)	2.23 (1.21, 4.10)*
Stressful life events	No	232	127	1	1
	Yes	17	35	3.76(2.02 6.98)	3.10 (1.60, 6.01)*
History of still birth	No	231	138	1	1
	Yes	18	24	2.23(1.16, 4.26)	1.32(0.60, 2.87)

COR Crude Odds Ratio, AOR Adjusted Odds Ratio, *P < 0.05

of sleep when compared with participants with good sleep hygiene practice. The significant association between poor sleep hygiene practice and poor sleep quality in the present study is similar to previous Ethiopian studies [42, 49] and a study done in Nigeria [56]. The possible explanation might be the risk of poor sleep quality in study subjects who had poor sleep hygiene practice might be justified by poor knowledge regarding the importance of sleep hygiene practice leading to poor sleep quality [57, 58].

Finally, we found an association between comorbid general anxiety symptoms and poor sleep quality. This is probably due to the fact that individuals, who had general anxiety symptoms, have an impact on poor quality of life and lead to poor sleep quality [59]. The finding is similar to a study done in Ethiopia [41] and the US [60].

Limitation of the study

This study didn't show the cause and effect relationship of the dependent and determinant variables. The other limitations might be interviewer bias and recall problems. Finally, we have not assessed factors like chronic physical illness, perceived stigma due to HIV/AIDS, depression, HIV- status of the previous child, current use of substances (alcohol, Khat, and tobacco), and other comorbid mental illnesses that may have higher significance for poor sleep quality.

Conclusion

The overall magnitude of poor sleep quality among HIV-positive pregnant women was high. Stressful life events; poor sleep hygiene practice, unplanned pregnancy, and comorbid general anxiety symptoms were the determinant factors of poor sleep quality. Therefore; we recommend that health care providers working in PMTCT clinics focused on early regular screening for sleep disturbance, and give higher attention to HIV-positive pregnant women who had an unplanned pregnancy, poor sleep hygiene practice, stressful life events, and comorbid general anxiety symptoms. Strengthening the referral linkage between health care providers working in PMTCT clinics and mental health specialists should be promoted.

Abbreviations

PSQI: Pittsburgh Sleep Quality Index; GAD-7: General anxiety disorder (GAD-7); SHI: Sleep Hygiene Index; PMTCT: Prevention of mothers-to-child-transmission; HIV: Human immune deficiency virus; AIDS: Acquired Immune Deficiency Syndrome; LTE: List of Threatening Experiences.

Acknowledgements

We would like to acknowledge Debre Tabor University for supporting ethical clearance and to data collectors and supervisors. We have sincere thanks to study subjects for giving their time.

Authors' contributions

GL conceived the study supported by GM, SA, AB, and TD. All authors participated in the development of the overall study design. Finally, the authors were involved in drafting, finalizing, and approving the manuscript.

Funding

Not applicable.

Availability of data and materials

all data are available in the manuscript.

Declarations

Ethics approval and consent to participate

All methodological procedures were performed under the declaration of Helsinki. The ethical approval was obtained from the ethical review committee of Debre Tabor University. Informed consent was taken from the participants. Respondents who agreed to participate in the study gave written consent. Informed consent for illiterate study subjects was obtained from their legally authorized representative after awareness of the purpose, and benefits of the study.

Consent for publication

Not applicable.

Competing interests

The authors have declared that no competing interests.

Author details

¹Department of Psychiatry, College of Medicine and Health Sciences, Debre Tabor University, Debre Tabor, Ethiopia. ²Department of Pediatrics and Child Health Nursing, College of Health Sciences, Debre Tabor University, Debre Tabor, Ethiopia.

Received: 17 May 2022 Accepted: 17 August 2022

Published online: 19 August 2022

References

- Nelson KL, Davis JE, Corbett CF, editors. Sleep quality: An evolutionary concept analysis. *Nursing Forum*; 2022: Wiley Online Library.
- Edition F. Diagnostic and statistical manual of mental disorders. *Am Psychiatric Assoc.* 2013;21:591–643.
- UNAIDS U. Fact sheet-Latest statistics on the status of the AIDS epidemic. 2016. 2017.
- Global H. AIDS statistics—2018 fact sheet. Geneva: UNAIDS. 2019.
- Control CfD, Prevention. Insufficient sleep is a public health problem. 2015.
- Oshinaike O, Akinbami A, Ojelabi O, Dada A, Dosunmu A, John Olabode S. Quality of sleep in an HIV population on antiretroviral therapy at an urban tertiary centre in Lagos, Nigeria. *Neurol Res Int.* 2014;2014. <https://doi.org/10.1155/2014/298703>.
- Altevogt BM, Colten HR. Sleep disorders and sleep deprivation: an unmet public health problem. 2006.
- Stranges S, Tigbe W, Gómez-Olivé FX, Thorogood M, Kandala N-B. Sleep problems: an emerging global epidemic? Findings from the INDEPTH WHO-SAGE study among more than 40,000 older adults from 8 countries across Africa and Asia. *Sleep.* 2012;35(8):1173–81.
- Reid S, Dwyer J. Insomnia in HIV infection: a systematic review of prevalence, correlates, and management. *Psychosomatic Med.* 2005;67(2):260–9.
- Bisong E. Predictors of sleep disorders among HIV out-patients in a tertiary hospital. *Recent Adv Biol Med.* 2017;3(2017):2747.
- Irwin MR, Olmstead R, Carroll JE. Sleep disturbance, sleep duration, and inflammation: a systematic review and meta-analysis of cohort studies and experimental sleep deprivation. *Biol Psychiatry.* 2016;80(1):40–52.
- Laposky AD, Pemberton VL. Sleep-disordered breathing and pregnancy-related cardiovascular disease. *J Women's Health.* 2021;30(2):194–8.

13. Micheli K, Komninos I, Bagkeris E, Roumeliotaki T, Koutis A, Kogevinas M, et al. Sleep patterns in late pregnancy and risk of preterm birth and fetal growth restriction. *Epidemiol.* 2011;738–44.
14. Du M, Liu J, Han N, Zhao Z, Yang J, Xu X, et al. Maternal sleep quality during early pregnancy, risk factors and its impact on pregnancy outcomes: a prospective cohort study. *Sleep Med.* 2021;79:11–8.
15. Nodine PM, Matthews EE. Common sleep disorders: management strategies and pregnancy outcomes. *J Midwifery Women's Health.* 2013;58(4):368–77.
16. Naghi I, Keypour F, Ahari S, Tavalai S, Khak M. Sleep disturbance in late pregnancy and type and duration of labour. *J Obstetr Gynaecol.* 2011;31(6):489–91.
17. Warland J, Dorrian J, Morrison JL, O'Brien LM. Maternal sleep during pregnancy and poor fetal outcomes: a scoping review of the literature with meta-analysis. *Sleep Med Rev.* 2018;41:197–219.
18. Qiu C, Sanchez SE, Gelaye B, Enquobahrie DA, Ananth CV, Williams MA. Maternal sleep duration and complaints of vital exhaustion during pregnancy is associated with placental abruption. *J Maternal-Fetal Neonatal Med.* 2015;28(3):350–5.
19. Cai S, Tan S, Gluckman PD, Godfrey KM, Saw S-M, Teoh OH, et al. Sleep quality and nocturnal sleep duration in pregnancy and risk of gestational diabetes mellitus. *Sleep.* 2017;40(2):zsw058.
20. Sharma S, Nehra A, Sinha S, Soneja M, Sunesh K, Sreenivas V, et al. Sleep disorders in pregnancy and their association with pregnancy outcomes: a prospective observational study. *Sleep Breath.* 2016;20(1):87–93.
21. Pauley AM, Moore GA, Mama SK, Molenaar P, Symons Downs D. Associations between prenatal sleep and psychological health: a systematic review. *J Clin Sleep Med.* 2020;16(4):619–30.
22. Li R, Zhang J, Zhou R, Liu J, Dai Z, Liu D, et al. Sleep disturbances during pregnancy are associated with cesarean delivery and preterm birth. *J Maternal-Fetal Neonatal Med.* 2017;30(6):733–8.
23. Okun ML, Schetter CD, Glynn LM. Poor sleep quality is associated with preterm birth. *Sleep.* 2011;34(11):1493–8.
24. Wang L, Jin F. Association between maternal sleep duration and quality, and the risk of preterm birth: a systematic review and meta-analysis of observational studies. *BMC Pregnancy Childbirth.* 2020;20(1):1–13.
25. Okun ML, Hanusa BH, Hall M, Wisner KL. Sleep complaints in late pregnancy and the recurrence of postpartum depression. *Behav Sleep Med.* 2009;7(2):106–17.
26. Dorheim SK, Bjorvatn B, Eberhard-Gran M. Can insomnia in pregnancy predict postpartum depression? A longitudinal, population-based study. *PLoS one.* 2014;9(4):e94674.
27. Pietikäinen JT, Polo-Kantola P, Pölkki P, Saarenpää-Heikkilä O, Paunio T, Paaonon EJ. Sleeping problems during pregnancy—a risk factor for postnatal depressiveness. *Arch Women's Mental Health.* 2019;22(3):327–37.
28. Lee J, Eklund EE, Lambert-Messerlian G, Palomaki GE, Butterfield K, Curran P, et al. Serum progesterone levels in pregnant women with obstructive sleep apnea: a case control study. *J Women's Health.* 2017;26(3):259–65.
29. Xu X, Liu D, Zhang Z, Sharma M, Zhao Y. Sleep duration and quality in pregnant women: a cross-sectional survey in China. *Int J Environ Res Public Health.* 2017;14(7):817.
30. Yang Y, Li W, Ma T-J, Zhang L, Hall BJ, Ungvari GS, et al. Prevalence of poor sleep quality in perinatal and postnatal women: a comprehensive meta-analysis of observational studies. *Front Psychiatry.* 2020;11:161.
31. Liu R-Q, Bloom MS, Wu Q-Z, He Z-Z, Qian Z, Stamatakis KA, et al. Association between depressive symptoms and poor sleep quality among Han and Manchu ethnicities in a large, rural, Chinese population. *PLoS one.* 2019;14(12):e0226562.
32. Christian LM, Carroll JE, Porter K, Hall MH. Sleep quality across pregnancy and postpartum: effects of parity and race. *Sleep Health.* 2019;5(4):327–34.
33. Huong NTT, Thuy NTH. Quality of sleep among pregnant women. *Int J Clin Med.* 2018;10(1):16–25.
34. Einstein A, Podolsky B, Rosen N. Can quantum-mechanical description of physical reality be considered complete? *Phys Rev.* 1935;47:777–80.
35. Teong ACA, Diong AX, Omar SZ, Tan PC. The impact of self-reported sleep on caesarean delivery in women undergoing induction of labour: a prospective study. *Scientific reports.* 2017;7(1):1–8.
36. Khadka R, Hong SA, Chang Y-S. Prevalence and determinants of poor sleep quality and depression among postpartum women: a community-based study in Ramechhap district Nepal. *Int Health.* 2020;12(2):125–31.
37. Ko SH, Chang SC, Chen CH. A comparative study of sleep quality between pregnant and nonpregnant Taiwanese women. *J Nurs Scholarship.* 2010;42(1):23–30.
38. Gelaye B, Addae G, Neway B, Larrabure-Torrevalva GT, Qiu C, Stoner L, et al. Poor sleep quality, antepartum depression and suicidal ideation among pregnant women. *J Affect Disord.* 2017;209:195–200.
39. Brand S, Kirov R. Sleep and its importance in adolescence and in common adolescent somatic and psychiatric conditions. *Int J General Med.* 2011;4:425.
40. Daubert E, French AL, Burgess HJ, Sharma A, Gustafson D, Cribbs SK, et al. Association of Poor Sleep With Depressive and Anxiety Symptoms by HIV Disease Status: Women's Interagency HIV Study. *J Acquired Immune Deficiency Syndromes (1999).* 2022;89(2):222.
41. Dule A, Abdu Z, Hajure M, Mohammedhusein M. Sleep Quality Among Pregnant Women Amidst COVID-19: Association with Mental Wellbeing and Self-efficacy. *Am J Health Res.* 2021;9(6):238–45.
42. Getachew Y, Azale T, Necho M. Poor sleep quality and associated factors among prisoners of the Diredawa correctional facility in eastern Ethiopia. *Ann General Psychiatry.* 2020;19(1):1–9.
43. Jemere T, Getahun B, Tadele F, Kefale B, Walle G. Poor sleep quality and its associated factors among pregnant women in Northern Ethiopia, 2020: A cross sectional study. *PLoS One.* 2021;16(5):e0250985.
44. Anbesaw T, Abebe H, Kassaw C, Bete T, Molla A. Sleep quality and associated factors among pregnant women attending antenatal care at Jimma Medical Center, Jimma, Southwest Ethiopia, 2020: cross-sectional study. *BMC Psychiatry.* 2021;21(1):1–11.
45. Amare NS, Chekol B, Aemro A. Determinants of Poor Sleep Quality During the COVID-19 Pandemic Among Women Attending Antenatal Care Services at the Health Facilities of Debre Berhan Town, Ethiopia: An Institutional-Based Cross-Sectional Study. *Front Psychiatry.* 2022;13. <https://doi.org/10.3389/fpsy.2022.841097>.
46. Li Y, Gu S, Wang Z, Li H, Xu X, Zhu H, et al. Relationship between stressful life events and sleep quality: rumination as a mediator and resilience as a moderator. *Front Psychiatry.* 2019;10:348.
47. Li Y-L, Qiu D, Hu C, Ouyang F-Y, He J, Zang D-F, et al. Stressful life events and poor sleep quality: a cross-sectional survey in the Chinese governmental employees. *Sleep Med.* 2021;85:123–30.
48. Ju YJ, Lee JE, Choi D-W, Han K-T, Lee SY. Association between perceived environmental pollution and poor sleep quality: results from nationwide general population sample of 162,797 people. *Sleep Med.* 2021;80:236–43.
49. Mengistu N, Azale T, Yimer S, Fikreyesus M, Melaku E, Shumye S. Quality of sleep and associated factors among people living with HIV/AIDS on follow up at Ethiopian Zewditu Memorial Hospital, 2018. *Sleep Sci Pract.* 2021;5(1):1–8.
50. Salahuddin M, Maru TT, Kumalo A, Pandi-Perumal SR, Bahammam AS, Manzar MD. Validation of the Pittsburgh sleep quality index in community dwelling Ethiopian adults. *Health Qual Life Outcomes.* 2017;15(1):1–7.
51. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Internal Med.* 2006;166(10):1092–7.
52. Cho S, Kim G-S, Lee J-H. Psychometric evaluation of the sleep hygiene index: a sample of patients with chronic pain. *Health Qual Life Outcomes.* 2013;11(1):1–7.
53. Brugha T, Bebbington P, Tennant C, Hurry J. The List of Threatening Experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychol Med.* 1985;15(1):189–94.
54. Baddam SK, Olvera RL, Canapari CA, Crowley MJ, Williamson DE. Childhood trauma and stressful life events are independently associated with sleep disturbances in adolescents. *Behav Sci.* 2019;9(10):108.
55. Richardson GS. Human physiological models of insomnia. *Sleep Med.* 2007;8:S9–14.
56. Fakorede O, Onifade P, Majekodunmi O. Sleep efficiency and sleep quality among male inmates in Nigeria. *Sleep Med.* 2015(16):S58. <https://doi.org/10.1016/j.sleep.2015.02.146>.
57. Brown FC, Buboltz WC Jr, Soper B. Development and evaluation of the Sleep Treatment and Education Program for Students (STEPS). *J Am Coll Health.* 2006;54(4):231–7.

58. O'Quinn S. Inadequate sleep hygiene practices: level of community awareness. *Can J Res Ther.* 2003;9:21–9.
59. Fu W, Wang C, Zou L, Guo Y, Lu Z, Yan S, et al. Psychological health, sleep quality, and coping styles to stress facing the COVID-19 in Wuhan, China. *Transl Psychiatry.* 2020;10(1):1–9.
60. Daubert E, French AL, Burgess HJ, Sharma A, Gustafson D, Cribbs SK, et al. Association of Poor Sleep With Depressive and Anxiety Symptoms by HIV Disease Status: Women's Interagency HIV Study. *J Acquired Immune Deficiency Syndromes.* 2022;89(2):222–30.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

