

RESEARCH ARTICLE

Open Access

The association between parental history of diagnosed mood/anxiety disorders and psychiatric symptoms and disorders in young adult offspring

Nancy CP Low^{1*}, Erika Dugas², Evelyn Constantin³, Igor Karp^{2,4}, Daniel Rodriguez⁵ and Jennifer O'Loughlin^{2,4,6}

Abstract

Background: Parental history of mood or anxiety disorders is one of the strongest and most consistent risk factors for the development of these disorders in offspring. Gaps remain however in our knowledge of whether maternal or paternal disorders are more strongly associated with offspring disorders, and whether the association exists in non-clinical samples. This study uses a large population-based sample to test if maternal or paternal history of mood and/or anxiety disorders increases the risk of mood and/or anxiety disorders, or symptoms of specific anxiety disorders, in offspring.

Methods: Data were drawn from the Nicotine Dependence in Teens Study, a prospective cohort investigation of 1293 grade 7 students. Data on mental health outcomes were collected in mailed self-report questionnaires when participants were aged 20.4 (0.7) years on average. Parental data were collected in mailed self-report questionnaires. This current analysis pertains to 564 participants with maternal and/or paternal data. The association between maternal and paternal history and each of diagnosed anxiety disorder, diagnosed mood disorder, and symptoms of specific anxiety disorders in offspring was studied in multivariate logistic regression.

Results: A higher proportion of mothers than fathers had a diagnosed mood/anxiety disorder (23% versus 12%). Similarly, 14% of female offspring had a diagnosed mood/anxiety disorder, compared to 6% of male offspring. The adjusted odds ratio (95% confidence interval) for maternal history was 2.2 (1.1, 4.5) for diagnosed mood disorders, 4.0 (2.1, 7.8) for diagnosed anxiety disorders, and 2.2 (1.2, 4.0) for social phobia symptoms. Paternal history was not associated with any of the mental health outcomes in offspring.

Conclusion: Maternal, but not paternal mood/anxiety disorders were associated with diagnosed psychiatric disorders, as well as symptoms of specific anxiety disorders, in offspring. Efforts to detect mood and anxiety disorders in offspring with a maternal history should be encouraged.

Keywords: Familial risk, Parental history, Offspring, Psychiatric, Mood, Anxiety, Panic

* Correspondence: nancy.low@mcgill.ca

¹Department of Psychiatry, McGill University, 1033 Pine Avenue West, Montréal, QC H3A 1A1, Canada

Full list of author information is available at the end of the article

Background

Mood and anxiety disorders are chronic conditions that affect up to 15-20% of adolescents and young adults [1-3]. These disorders are associated with academic failure [4-6], difficulties finding and sustaining employment [7,8], problems in relationships with family, friends and romantic partners [6,9-12], substance misuse [13-15], and suicide [16-18]. Although onset of symptoms of mood and anxiety disorders often occurs during childhood or adolescence [2,19,20] diagnosis and treatment may not occur until up to 14 years later for mood disorders, and 30 years later for anxiety disorders [21]. The years between first symptoms and diagnosis represent a critical period for preventive intervention and treatment which could reduce the future burden of mood and anxiety disorders.

Parental history of mood and anxiety disorders is one of the strongest and most consistent risk factors for the development of these disorders in offspring [22-24]. Based on findings from clinical samples (i.e., in which either parents or offspring are recruited in treatment settings), the increased risk to offspring with a parental history relative to offspring with no parental history, ranges from 2 to 4-fold for depression [25-34] and from 2 to 7-fold for anxiety [29,31,32,35-37] including panic [34,38], social phobia [38] and generalized anxiety disorders [34,38,39]. The increase in risk is greater for maternal versus paternal history [27-30] although not all studies report differences [26,28]. In addition to inconsistent findings across studies, most studies that use clinical samples are limited by small sample sizes, examination of one parent only (predominantly the mother), or assessment of offspring who may not yet be old enough to manifest illness.

In contrast to studies of clinical samples, reports based on population-based samples in which self-report or direct assessments of psychiatric disorders are collected from both parents and offspring are rare. Studies that do exist [40-44] report a range in risk from no elevated risk to an increase in risk of 1.5 to 2.5-fold. Again, inconsistencies are reported in terms of maternal versus paternal history. Two studies suggest maternal but not paternal associations, [40,42] and two suggest both [41,44]. To date, no population-based studies describe the risk of parent self-reported psychiatric history on specific offspring anxiety disorders (i.e. panic, social phobia and generalized anxiety disorders). Thus, it remains unclear whether the association between parental history and offspring psychiatric disorders exists in population-based samples of youth.

The objective of this study is to test if maternal or paternal history of diagnosed mood or anxiety disorders is associated with the risk of mood or anxiety disorders, or with symptoms of specific anxiety disorders in a large population-based cohort.

Methods

Data were drawn from the Nicotine Dependence in Teens (NDIT) Study, a prospective cohort investigation of 1293 students recruited in 1999–2000 from all grade 7 classes in a convenience sample of 10 secondary schools in Montreal, Canada [45]. Self-report questionnaires were administered in classrooms every 3 months during secondary school from grade 7–11, for a total of 20 survey cycles during the five years of secondary school. In 2007–8, when participants were aged 20.4 (0.7) years on average, data were collected in mailed self-report questionnaires from a total of 880 participants (68% of 1293 participants) in survey cycle 21. In addition, in 2009–10, data were collected from parents in three mailed self-report questionnaires including a questionnaire for mothers (n=597; 97.7% biological mothers), a questionnaire for fathers (n=478; 97.1% biological fathers), and a questionnaire which was completed by either parent that collected data about the NDIT participant (n =647).

This current analysis pertains to NDIT participants (herein referred to as offspring) who completed survey cycle 21 and whose parents completed the self-report parental questionnaires. Two datasets were constructed, one that consisted of offspring and maternal self-report data, and the second that consisted of offspring and paternal self-report data (referred to herein as the “mother database” and “father database”, respectively). The mother dataset consisted of 564 (64% of 880 eligible offspring) participants with complete data, and the father dataset consisted of 454 (52% of 880 eligible offspring) participants with complete data. All offspring and parents provided informed consent. The study received ethics approval from the Montreal Department of Public Health Ethics Review Committee, the McGill University Faculty of Medicine Institutional Review Board, and the Ethics Research Committee of the Centre de Recherche du Centre Hospitalier de l’Université de Montréal.

Study variables

Socio-demographic data pertaining to offspring included age, sex, language spoken most often at home (French, other), mother’s level of education (attended university, did not attend university), annual household income (<\$30,000, \$30,000-\$99,999, ≥\$100,000), and born in Canada (yes, no).

Data on diagnosed mental health disorders (yes, no) among mothers and fathers were collected in the parent questionnaires by: “Have you ever been diagnosed by a health professional with any of the following...”: (i) anxiety disorder (phobia, obsessive-compulsive disorder, panic attacks, generalized anxiety disorder (GAD)); (ii) depression; (iii) bipolar disorder; and (iv) postpartum depression (for mothers only). Data on age at diagnosis

were collected if the parent endorsed being diagnosed by a health professional. For multivariable analyses, anxiety disorders, depression, bipolar disorder and postpartum depression were grouped into a single variable termed “anxiety or mood disorder”.

Diagnosed mental health disorders (yes, no) among offspring were measured in the mailed self-report questionnaire in survey cycle 21 by: “Has a health professional ever diagnosed you with the following...”: (i) mood disorder (depression, bipolar disorder); (ii) anxiety disorder (phobia, fear of social situations, obsessive-compulsive disorder, panic disorder, GAD). Data on age at diagnosis were collected if the offspring endorsed being diagnosed by a health professional.

Lifetime anxiety symptoms of panic disorder, GAD, and social phobia among offspring were assessed in survey cycle 21 using the Composite International Diagnostic Interview (CIDI) screening questions [46]. Symptoms of panic disorder (yes, no) were assessed by “Have you ever had any of the following”: (i) attack of fear or panic when all of sudden you felt very frightened, anxious or uneasy; and (ii) attack when all of sudden, you became dizzy, very uncomfortable, short of breath, dizzy, nauseous, your heart pounded, or you thought that you might lose control, die or go crazy. Symptoms of GAD (yes, no) were assessed by “Have you ever had any of the following”: (i) a time when you were a “worrier” (when you worried a lot more about things than other people with the same problem); (ii) a period lasting 6 month or longer when you were anxious or worried most days?; (iii) a time when you were much more nervous or anxious than most other people with the same problems. Symptoms of social phobia (yes, no) were assessed by “Have you ever had any of the following”: (i) a time when you felt very afraid or really shy meeting new people, going to parties, going on a date; and (ii) a time when you felt very afraid or uncomfortable when you had to do something in front of a group of people (giving a speech, speaking in class). Data on age at onset of symptom(s) were collected if the offspring endorsed ever having the symptom(s). For each anxiety subtype, offspring who responded “yes” to at least one symptom within each anxiety subtype were categorized as screening positively for that anxiety subtype. The sensitivity and specificity of the CIDI anxiety disorder screening questions compared to full diagnoses generated by the administration of the complete CIDI instrument range from 89-95% and 51-98%, respectively [47].

Data analysis

Differences in socio-demographic characteristics between participants retained and not retained for analysis were tested in two-sample t-tests for continuous variables, and in chi-square tests for binary variables.

Because early analyses suggested that there was no statistically significant interaction between offspring sex and either maternal or paternal history of mood and/or anxiety disorders, all multivariate analyses were pooled by sex. The independent association between each of maternal history and paternal history and each of the five (binary) outcome variables (i.e., offspring diagnosed anxiety disorder, offspring diagnosed mood disorder, and offspring symptoms of panic disorder, social phobia, and GAD) was assessed in separate multivariate logistic regression models controlling for maternal education and offspring sex (i.e., the only socio-demographic variables associated with the mental health outcomes at $p \leq 0.25$ in univariate analyses (data not shown)). In secondary analyses, parental history of depression, bipolar and anxiety disorders were studied as separate exposures (Additional file 1: Table S1). In the paternal history models, we also tested for an interaction between fathers living with offspring (duration in years) and paternal history. All analyses were conducted using SAS software, Version 9.2.

Results

While several socio-demographic characteristics of offspring retained for analysis were statistically significantly different from those of offspring not retained, the differences were not substantively important with one exception (Table 1). In the father database, 49% of offspring retained for analysis had fathers who were university-educated, compared to 40% of those not retained.

Mothers and fathers were aged 52 (sd=7) and 54 (sd=7) years on average, respectively at the time they completed the self-report parental questionnaires. Compared to fathers, higher proportions of mothers had been diagnosed with both mood and anxiety disorders, with the exception of bipolar disorder (Table 2). Specifically, 23% of mothers had been diagnosed with either a mood or anxiety disorder, compared to 12% of fathers. Five percent of mothers had been diagnosed with both disorders, compared to 2% of fathers.

Similarly, statistically significantly higher proportions of female offspring (14%) had been diagnosed with either a mood or anxiety disorder, compared to male offspring (6%) (Table 3). Symptoms of social phobia were the most frequently endorsed anxiety subtype among both male and female offspring. With the exception of social phobia symptoms (for which there was no statistically significant sex difference), higher proportions of female than male offspring endorsed each anxiety subtype.

A diagnosis of mood and/or anxiety disorders in mothers was associated with a 2- and 4-fold increase in the odds of offspring having a diagnosed mood and anxiety disorder, respectively (Table 4). Maternal history was also statistically significantly associated with an increased likelihood of social phobia symptoms in offspring.

Table 1 Comparison of selected socio demographic characteristics of offspring retained for analysis with those of offspring not retained, NDI 2007-8

	Mother database			Father database		
	Retained (n = 564)	Not retained (n = 316)	p-value for difference	Retained (n = 454)	Not retained (n = 426)	p-value for difference
Age, y, mean (sd)	20.3 (0.7)	20.6 (0.8)	<0.0001	20.3 (0.7)	20.5 (0.8)	<0.0001
Male %	46.6	44.8	0.6604	47.0	44.7	0.5323
Mother university-educated, %	46.7	43.6	0.6654	49.2	39.5	0.0090
Household income, %						
<\$30,000	36.7	47.5	0.0057	35.9	45.4	0.0002
\$30,000-	44.7	41.5		42.9	44.5	
\$99,999	18.6	11.0		21.3	10.1	
≥\$100,000						
Canadian born, %	95.6	89.5	0.0009	95.8	90.8	0.0044
French-speaking,%	33.6	27.9	0.1774	28.4	33.6	0.1040

Paternal history was not statistically significantly associated with any of the mental health outcomes in offspring (Table 4). There were no statistically significant interaction between duration of time spent living with father (mean (sd) = 20 (4) years) and paternal mood and/or anxiety disorders.

Supplementary analyses (Additional file 1: Table S1) suggested that a depression diagnosis in mothers was associated with a 3-fold, 2-fold and 2-fold increase in the odds of offspring having diagnosed anxiety disorder, panic and GAD symptoms, respectively. Further, a diagnosis of anxiety disorder in mothers was associated with a 3-fold, 5-fold and 2-fold increase in the odds of offspring having diagnosed mood and anxiety disorders, and social phobia symptoms, respectively. Finally, a bipolar disorder diagnosis in fathers was associated with a 13-fold increase in the odds of offspring having diagnosed anxiety disorder. With respect to the temporal sequence of parental and offspring diagnoses and symptoms, maternal age of diagnosis of anxiety and mood

disorders (39 (sd=9) and 39 (sd=10) years, respectively) preceded the maternal age at which offspring were diagnosed with mood and anxiety disorders (54 (sd=7) and 56 (sd=7) years, respectively) and when offspring first had social phobia symptoms (53 (sd=7) years), GAD symptoms (53 (sd=6) years), and panic symptoms (53 (sd=6) years).

Similarly for fathers, paternal age of diagnosis of anxiety and mood disorders (44 (sd=12) and 42 (sd=11) years, respectively) preceded paternal age when offspring were diagnosed with mood and anxiety disorders (58 (sd=6) and 60 (sd=6) years, respectively) and when offspring first had social phobia symptoms (55 (sd=7) years), GAD symptoms (57 (sd=7) years), and panic symptoms (55 (sd= 8) years).

Discussion

Despite differences in methodologies, the main finding of this analysis - the strong and statistically significant association between maternal history and mood and anxiety disorders in offspring - is consistent with extant population-based studies [40-42,44,48]. The magnitude of the associations observed concurs with that reported in previous population-based and clinical studies [25-33,35,36,38,39]. In addition, maternal history was related to symptoms of social phobia in offspring.

These findings emphasize the importance of ongoing research on familial clustering of mood and anxiety disorders. Despite investigation of conceptually credible mechanisms of the maternal history-offspring disorder association, which include genetic transmission [49-51], parenting behavior [52,53], and depressogenic child-rearing environments [54-57] recent reviews suggest that these mechanisms explain little of the variation in the association [23,52,53]. This has led to the studies that investigate parental history, parenting behavior, and familial environments

Table 2 Proportion of parents diagnosed with a mood and/or anxiety disorder, NDI 2009-10

	Mother (n=564) %	Father (n=454) %
Mood disorder		
Depression	18.2	9.9
Bipolar	1.1	1.1
Postpartum depression	4.0	Not applicable
Any mood disorder	19.3	9.7
Anxiety disorder*	8.9	4.1
Mood and/or anxiety disorder		
Mood or anxiety disorder	23.0	11.8
Mood and anxiety disorder	4.8	1.6

*Includes phobias, obsessive compulsive disorder, panic attacks, and generalized anxiety disorder.

Table 3 Proportion of offspring diagnosed with mood and/or anxiety disorders, and proportion with symptoms of specific anxiety disorders, NDIT 2007–8

	Total (n= 564) %	Female (n=302) %	Male (n=262) %	p-value for difference by sex
Diagnosed				
Mood disorder	6.9	9.6	3.8	0.0116
Anxiety disorder	7.6	10.3	4.6	0.0180
Mood or anxiety disorder	10.3	13.9	6.1	0.0039
Mood and anxiety disorder	4.3	6.0	2.3	0.0530
Anxiety symptoms				
Panic disorder	15.7	21.6	8.8	<0.0001
Generalized anxiety disorder (GAD)	9.4	13.7	4.6	0.0004
Social phobia	24.0	26.5	21.2	0.1684

concurrently [54,55]. More recent studies discuss other factors that may explain more of the variance in the maternal history-offspring disorder association including mindful parenting [58-60], positive maternal behaviours during problem-solving tasks with children, use of an avoidance conflict-resolution approach [55], positive offspring coping skills [61] and familial environments characterized by critical expressed emotion [62].

In this study, we did not detect any association between paternal history and offspring disorder, which

supports several population-based studies [40,42], but not others [41,44]. This inconsistency may relate, at least in part, to methodological differences between studies. In contrast to several reports [40,42], our study required that parents be diagnosed by a health care professional. Men seek treatment for mental health disorders much less frequently than women [63] and therefore some fathers in our sample could have met the diagnostic criteria for a mood and/or anxiety disorder, but never sought or received treatment. Further, paternal history may be related to offspring outcomes that were not within the scope of our study, such as conduct disorder and attention-deficit hyperactivity disorder [42,44]. Detection of an association between paternal and offspring disorders may have been limited by low power [19,64,65]. Finally, paternal and offspring mood and/or anxiety disorders may not be related, at least in our sample.

Our findings suggest that offspring with maternal mood and anxiety disorder histories should be monitored for social phobia symptoms, in addition to mood and anxiety disorders. Adequate screening and treatment of parents, and in particular, mothers, is also essential as parents are instrumental in effective treatment of children and adolescents with mood and anxiety symptoms and disorders [66]. Mothers can ensure that children do their homework, practice skills taught during treatment sessions, and monitor symptoms and response to interventions. Concomitant treatment of mothers with children being treated in clinics has improved outcomes in offspring [67].

Table 4 Adjusted odds ratios (OR) and 95% confidence intervals (CI) for the association between diagnosed parental mood and/or anxiety disorders and mental health outcomes among offspring, NDIT 2007-10

	n	Offspring diagnosis		Anxiety symptoms in offspring		
		Mood disorder OR _{adj} (95% CI)	Anxiety disorder OR _{adj} (95% CI)	Social phobia OR _{adj} (95% CI)	Generalized anxiety disorder (GAD) OR _{adj} (95% CI)	Panic disorder OR _{adj} (95% CI)
Mother diagnosed with mood and/or anxiety disorder	564*	2.2 (1.1,4.5)	4.0 (2.1,7.8)	2.2 (1.2,4.0)	1.8 (0.7,4.1)	1.7 (0.8,3.5)
p-value		0.0263	<0.0001	0.0131	0.1966	0.1497
Offspring sex		2.4 (1.1,5.3)	2.4 (1.2,5.0)	1.4 (1.0,2.2)	3.6 (1.9,7.4)	2.6 (1.6,4.5)
p-value		0.0249	0.0183	0.0752	0.0002	0.0003
Mother university-educated (yes, no)		0.5 (0.2,1.0)	0.50 (0.2,1.00)	1.3 (0.9,2.0)	2.0 (1.1,3.6)	0.7 (0.4,1.2)
p-value		0.0542	0.0583	0.1614	0.0233	0.2050
Father diagnosed with mood and/or disorder**	454*	0.9 (0.2,2.8)	**	0.5 (0.2,1.1)	0.8 (0.2,2.2)	**
p-value		0.8862		0.1263	0.7112	
Offspring sex		3.6 (1.4,11.1)		1.7 (1.1,2.7)	5.9 (2.6,15.9)	
p-value		0.0115		0.0236	0.0001	
Mother university-educated (yes, no)		0.4 (0.2,0.9)		1.1 (0.7,1.8)	1.9 (1.0,3.8)	
p-value		0.0429		0.6206	0.0622	

* n varies because of missing data.

**Model did not converge and is therefore not reported.

Limitations of this analysis include that mood and anxiety diagnostic data were based on self-report rather than structured interviews. Misclassification of the disorders may have attenuated estimates of the association between parental history and offspring mental health disorders. Ascertainment of disorders using diagnosis by a mental health professional may not reflect all cases of disorder. However, we did not collect data on access to services or frequency of consultation. The use of what is, in essence, cross-sectional data limits causal inference. However, age of parental diagnoses preceded age of offspring diagnoses and onset of anxiety symptoms.

Conclusion

In this study, maternal, but not paternal, mood and/or anxiety disorders were associated with mood and anxiety disorders in offspring. Depression among adults is a chronic, recurring disorder and therefore vigilance about its potential impact on offspring needs to be maintained. Clinically, it is critical when treating adults with depression to screen for symptoms in children, and similarly to screen for maternal disorders when treating children. Effective treatment of both offspring and parental disorders has been demonstrated, with optimal outcomes occurring when both are treated [68]. Future research should focus on ascertaining the precise timing and nature of the maternal symptoms that have the greatest negative impact on offspring. Further delineation of the natural course of depressive and anxiety disorders in high-risk offspring may help determine when and which symptoms first appear, which may lead to more timely intervention in both individuals and families.

Additional file

Additional file 1: Table S1. Adjusted odds ratios (OR) and 95% confidence intervals (CI) for the association between diagnosed parental mood and/or anxiety disorders and mental health outcomes among offspring, NDIT 2007–10.

Competing interests

The authors declare no competing interests.

Authors' contributions

Contribution of authors: NL defined the study question, reviewed the literature, contributed to the design of the analysis and interpretation of data, and wrote sections of the article. ED, EC, DR, IK reviewed the literature, contributed to the design of the analysis and interpretation of data, and wrote sections of the article. DR contributed to data analysis and drafted the data analysis section in the article. JOL designed the study, obtained the funding, developed the survey instruments, supervised data collection, contributed to the design of the analysis and interpretation of data, coordinated drafting the article, and critically reviewed and edited sections of the article. All authors reviewed the article and approved the final version.

Author details

¹Department of Psychiatry, McGill University, 1033 Pine Avenue West, Montréal, QC H3A 1A1, Canada. ²Centre de Recherche de Centre Hospitalier de l'Université de Montréal, 3875 Saint Urbain, Montréal, QC H2W 1V1,

Canada. ³Department of Pediatrics, McGill University, Montreal Children's Hospital, 2300 Tupper Street, Room C-538E, Montréal, QC H3H 1P3, Canada. ⁴Department of Social and Preventive Medicine, University of Montréal, 7171 Parc Avenue, Montréal, QC, Canada. ⁵Centers for Behavioral and Preventive Medicine, Department of Psychiatry and Human Behavior, Alpert Medical School, Brown University, The Miriam Hospital, Coro West, Suite 309, 164 Summit Avenue, Providence, RI 02906, USA. ⁶Institut national de santé publique du Québec, 190 Crémazie Blvd. East, Montréal, QC H2P 1E2, Canada.

Received: 17 February 2012 Accepted: 27 October 2012

Published: 5 November 2012

References

- Lewinsohn PM, Hops H, Roberts RE, Seeley JR, Andrews JA: **Adolescent psychopathology: I Prevalence and incidence of depression and other DSM-III-R disorders in high school students.** *J Abnorm Psychol* 1993, **102**(1):133–144.
- Costello EJ, Pine DS, Hammen C, March JS, Plotsky PM, Weissman MM, Biederman J, Goldsmith HH, Kaufman J, Lewinsohn PM, et al: **Development and natural history of mood disorders.** *Biol Psychiatry* 2002, **52**(6):529–542.
- Kashani JH, Orvaschel H: **A community study of anxiety in children and adolescents.** *Am J Psychiatry* 1990, **147**(3):313–318.
- Mazzone L, Ducci F, Scoto MC, Passaniti E, D'Arrigo VG, Vitiello B: **The role of anxiety symptoms in school performance in a community sample of children and adolescents.** *BMC Publ Health* 2007, **7**:347.
- Egger HL, Costello EJ, Angold A: **School refusal and psychiatric disorders: a community study.** *J Am Acad Child Adolesc Psychiatry* 2003, **42**(7):797–807.
- Jaycox LH, Stein BD, Paddock S, Miles JN, Chandra A, Meredith LS, Tanielian T, Hickey S, Burnam MA: **Impact of teen depression on academic, social, and physical functioning.** *Pediatrics* 2009, **124**(4):e596–e605.
- Lauber C, Bowen JL: **Low mood and employment in a community sample of individuals with affective disorders: a UK perspective.** *Int Rev Psychiatry* 2010, **22**(2):173–182.
- Cowell AJ, Luo Z, Masuda YJ: **Psychiatric disorders and the labor market: an analysis by disorder profiles.** *J Ment Health Policy Econ* 2009, **12**(1):3–17.
- Marazziti D, Dell'osso B, Catena Dell'Osso M, Consoli G, Del Debbio A, Mungai F, Vivarelli L, Albanese F, Piccinni A, Rucci P, et al: **Romantic attachment in patients with mood and anxiety disorders.** *CNS Spectr* 2007, **12**(10):751–756.
- Benvenuti A, Rucci P, Calugi S, Cassano GB, Miniati M, Frank E: **Relationship of residual mood and panic-agoraphobic spectrum phenomenology to quality of life and functional impairment in patients with major depression.** *Int Clin Psychopharmacol* 2010, **25**(2):68–74.
- Goppoldova E, Dragomirecka E, Motlova L, Hajek T: **Subjective quality of life in psychiatric patients: diagnosis and illness-specific profiles.** *Can J Psychiatry* 2008, **53**(9):587–593.
- Olatunji BO, Cisler JM, Tolin DF: **Quality of life in the anxiety disorders: a meta-analytic review.** *Clin Psychol Rev* 2007, **27**(5):572–581.
- Frojd S, Ranta K, Kaltiala-Heino R, Marttunen M: **Associations of social phobia and general anxiety with alcohol and drug use in a community sample of adolescents.** *Alcohol Alcohol* 2011, **46**(2):192–199.
- Wu P, Goodwin RD, Fuller C, Liu X, Comer JS, Cohen P, Hoven CW: **The relationship between anxiety disorders and substance use among adolescents in the community: specificity and gender differences.** *J Youth Adolesc* 2010, **39**(2):177–188.
- O'Neil KA, Conner BT, Kendall PC: **Internalizing disorders and substance use disorders in youth: comorbidity, risk, temporal order, and implications for intervention.** *Clin Psychol Rev* 2011, **31**(1):104–112.
- Greydanus DE, Calles J Jr: **Suicide in children and adolescents.** *Prim Care* 2007, **34**(2):259–273. abstract vi.
- Boden JM, Fergusson DM, Horwood LJ: **Anxiety disorders and suicidal behaviours in adolescence and young adulthood: findings from a longitudinal study.** *Psychol Med* 2007, **37**(3):431–440.
- Pickles A, Aglan A, Collishaw S, Messer J, Rutter M, Maughan B: **Predictors of suicidality across the life span: the Isle of Wight study.** *Psychol Med* 2010, **40**(9):1453–1466.
- Kessler RC, Ruscio AM, Shear K, Wittchen HU: **Epidemiology of anxiety disorders.** *Curr Top Behav Neurosci* 2010, **2**:21–35.

20. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE: **Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication.** *Arch Gen Psychiatry* 2005, **62**(6):593–602.
21. Wang PS, Angermeyer M, Borges G, Bruffaerts R, Tat Chiu W, DEG G, Fayyad J, Gureje O, Haro JM, Huang Y, et al: **Delay and failure in treatment seeking after first onset of mental disorders in the World Health Organization's World Mental Health Survey Initiative.** *World Psychiatry* 2007, **6**(3):177–185.
22. Beardslee WR, Versage EM, Gladstone TR: **Children of affectively ill parents: a review of the past 10 years.** *J Am Acad Child Adolesc Psychiatry* 1998, **37**(11):1134–1141.
23. Goodman SH, Rouse MH, Connell AM, Broth MR, Hall CM, Heyward D: **Maternal depression and child psychopathology: a meta-analytic review.** *Clin Child Fam Psychol Rev* 2011, **14**(1):1–27.
24. Ramchandani P, Psychogiou L: **Paternal psychiatric disorders and children's psychosocial development.** *Lancet* 2009, **374**(9690):646–653.
25. Weissman MM, John K, Merikangas KR, Prusoff BA, Wickramaratne P, Gammon GD, Angold A, Warner V: **Depressed parents and their children. General health, social, and psychiatric problems.** *Am J Dis Child* 1986, **140**(8):801–805.
26. Keller MB, Beardslee WR, Dorer DJ, Lavori PW, Samuelson H, Klerman GR: **Impact of severity and chronicity of parental affective illness on adaptive functioning and psychopathology in children.** *Arch Gen Psychiatry* 1986, **43**(10):930–937.
27. Lavori PW, Keller MB, Beardslee WR, Dorer DJ: **Affective disorder in childhood: separating the familial component of risk from individual characteristics of children.** *J Affect Disord* 1988, **15**(3):303–311.
28. Mitchell J, McCauley E, Burke P, Calderon R, Schloredt K: **Psychopathology in parents of depressed children and adolescents.** *J Am Acad Child Adolesc Psychiatry* 1989, **28**(3):352–357.
29. Weller RA, Kapadia P, Weller EB, Fristad M, Lazaroff LB, Preskorn SH: **Psychopathology in families of children with major depressive disorders.** *J Affect Disord* 1994, **31**(4):247–252.
30. Welner Z, Rice J: **School-aged children of depressed parents: a blind and controlled study.** *J Affect Disord* 1988, **15**(3):291–302.
31. Swartz HA, Shear MK, Wren FJ, Greeno CG, Sales E, Sullivan BK, Ludewig DP: **Depression and anxiety among mothers who bring their children to a pediatric mental health clinic.** *Psychiatr Serv* 2005, **56**(9):1077–1083.
32. Warner V, Mufson L, Weissman MM: **Offspring at high and low risk for depression and anxiety: mechanisms of psychiatric disorder.** *J Am Acad Child Adolesc Psychiatry* 1995, **34**(6):786–797.
33. Mufson L, Weissman MM, Warner V: **Depression and anxiety in parents and children: a direct interview study.** *J Anxiety Disord* 1992, **6**:1–13.
34. Biederman J, Faraone SV, Hirshfeld-Becker DR, Friedman D, Robin JA, Rosenbaum JF: **Patterns of psychopathology and dysfunction in high-risk children of parents with panic disorder and major depression.** *Am J Psychiatry* 2001, **158**(1):49–57.
35. Merikangas KR, Dierker LC, Szatmari P: **Psychopathology among offspring of parents with substance abuse and/or anxiety disorders: a high-risk study.** *J Child Psychol Psychiatry* 1998, **39**(5):711–720.
36. Turner SM, Beidel DC, Costello A: **Psychopathology in the offspring of anxiety disorders patients.** *J Consult Clin Psychol* 1987, **55**(2):229–235.
37. Beidel DC, Turner SM: **At risk for anxiety: I Psychopathology in the offspring of anxious parents.** *J Am Acad Child Adolesc Psychiatry* 1997, **36**(7):918–924.
38. Cooper PJ, Fearn V, Willetts L, Seabrook H, Parkinson M: **Affective disorder in the parents of a clinic sample of children with anxiety disorders.** *J Affect Disord* 2006, **93**(1–3):205–212.
39. Last CG, Hersen M, Kazdin AE, Francis G, Grubb HJ: **Psychiatric illness in the mothers of anxious children.** *Am J Psychiatry* 1987, **144**(12):1580–1583.
40. McClure EB, Brennan PA, Hammen C, Le Brocque RM: **Parental anxiety disorders, child anxiety disorders, and the perceived parent-child relationship in an Australian high-risk sample.** *J Abnorm Child Psychol* 2001, **29**(1):1–10.
41. Klein DN, Lewinsohn PM, Rohde P, Seeley JR, Olino TM: **Psychopathology in the adolescent and young adult offspring of a community sample of mothers and fathers with major depression.** *Psychol Med* 2005, **35**(3):353–365.
42. Brennan PA, Hammen C, Katz AR, Le Brocque RM: **Maternal depression, paternal psychopathology, and adolescent diagnostic outcomes.** *J Consult Clin Psychol* 2002, **70**(5):1075–1085.
43. Ramchandani PG, Stein A, O'Connor TG, Heron J, Murray L, Evans J: **Depression in men in the postnatal period and later child psychopathology: a population cohort study.** *J Am Acad Child Adolesc Psychiatry* 2008, **47**(4):390–398.
44. Renk K, Oliveros A, Roddenberry A, Klein J, Sieger K, Roberts R, Phares V: **The relationship between maternal and paternal psychological symptoms and ratings of adolescent functioning.** *J Adolesc* 2007, **30**(3):467–485.
45. O'Loughlin J, Karp I, Koulis T, Paradis G, Difranza J: **Determinants of first puff and daily cigarette smoking in adolescents.** *Am J Epidemiol* 2009, **170**(5):585–597.
46. Wittchen HU: **Reliability and validity studies of the WHO-Composite International Diagnostic Interview (CIDI): a critical review.** *J Psychiatr Res* 1994, **28**(1):57–84.
47. Wittchen HU, Boyer P: **Screening for anxiety disorders. Sensitivity and specificity of the Anxiety Screening Questionnaire (ASQ-15).** *Br J Psychiatry Suppl* 1998, **34**:10–17.
48. van Meurs I, Reef J, Verhulst FC, van der Ende J: **Intergenerational transmission of child problem behaviors: a longitudinal, population-based study.** *J Am Acad Child Adolesc Psychiatry* 2009, **48**(2):138–145.
49. Merikangas KR, Chakravarti A, Moldin SO, Araj H, Blangero JC, Burmeister M, Crabbe J Jr, Depaulo JR Jr, Foulks E, Freimer NB, et al: **Future of genetics of mood disorders research.** *Biol Psychiatry* 2002, **52**(6):457–477.
50. Lau JY, Eley TC: **The genetics of mood disorders.** *Annu Rev Clin Psychol* 2010, **6**:313–337.
51. Smoller JW, Block SR, Young MM: **Genetics of anxiety disorders: the complex road from DSM to DNA.** *Depress Anxiety* 2009, **26**(11):965–975.
52. McLeod BD, Weisz JR, Wood JJ: **Examining the association between parenting and childhood depression: a meta-analysis.** *Clin Psychol Rev* 2007, **27**(8):986–1003.
53. McLeod BD, Wood JJ, Weisz JR: **Examining the association between parenting and childhood anxiety: a meta-analysis.** *Clin Psychol Rev* 2007, **27**(2):155–172.
54. Foster CJ, Garber J, Durlak JA: **Current and past maternal depression, maternal interaction behaviors, and children's externalizing and internalizing symptoms.** *J Abnorm Child Psychol* 2008, **36**(4):527–537.
55. Marchand JF, Hock E: **Mothers' and fathers' depressive symptoms and conflict-resolution strategies in the marriage and children's externalizing and internalizing behaviors.** *J Genet Psychol* 2003, **164**(2):227–239.
56. Du Rocher Schudlich TD, Cummings EM: **Parental dysphoria and children's internalizing symptoms: marital conflict styles as mediators of risk.** *Child Dev* 2003, **74**(6):1663–1681.
57. Burke L: **The impact of maternal depression on familial relationships.** *Int Rev Psychiatry* 2003, **15**(3):243–255.
58. Parent J, Garai E, Forehand R, Roland E, Potts J, Haker K, Champion JE, Compas BE: **Parent Mindfulness and Child Outcome: The Roles of Parent Depressive Symptoms and Parenting.** *Mindfulness (N Y)* 2011, **1**(4):254–264.
59. Coatsworth JD, Duncan LG, Greenberg MT, Nix RL: **Changing parent's mindfulness, child management skills and relationship quality with their youth: Results from a randomized pilot intervention trial.** *J Child Fam Stud* 2010, **19**:203–217.
60. Duncan LG, Coatsworth JD, Greenberg MT: **A model of mindful parenting: implications for parent-child relationships and prevention research.** *Clin Child Fam Psychol Rev* 2009, **12**(3):255–270.
61. Compas BE, Champion JE, Forehand R, Cole DA, Reeslund KL, Fear J, Hardcastle EJ, Keller G, Rakow A, Garai E, et al: **Coping and parenting: Mediators of 12-month outcomes of a family group cognitive-behavioral preventive intervention with families of depressed parents.** *J Consult Clin Psychol* 2010, **78**(5):623–634.
62. Gibb BE, Uhrlass DJ, Grassia M, Benas JS, McGeary J: **Children's inferential styles, 5-HTTLPR genotype, and maternal expressed emotion-criticism: An integrated model for the intergenerational transmission of depression.** *J Abnorm Psychol* 2009, **118**(4):734–745.
63. Galdas PM, Cheater F, Marshall P: **Men and health help-seeking behaviour: literature review.** *J Adv Nurs* 2005, **49**(6):616–623.
64. Vesga-Lopez O, Schneier FR, Wang S, Heimberg RG, Liu SM, Hasin DS, Blanco C: **Gender differences in generalized anxiety disorder: Results from the national epidemiologic survey on alcohol and related conditions (NESARC).** *J Clin Psychiatry* 2008, **69**(10):1606–1616.

65. Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, Rush AJ, Walters EE, Wang PS: **The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R).** *JAMA* 2003, **289**(23):3095–3105.
66. Silverman WK, Ortiz CD, Viswesvaran C, Burns BJ, Kolko DJ, Putnam FW, Amaya-Jackson L: **Evidence-based psychosocial treatments for children and adolescents exposed to traumatic events.** *J Clin Child Adolesc Psychol* 2008, **37**(1):156–183.
67. Swartz HA, Frank E, Zuckoff A, Cyranowski JM, Houck PR, Cheng Y, Fleming MA, Grote NK, Brent DA, Shear MK: **Brief interpersonal psychotherapy for depressed mothers whose children are receiving psychiatric treatment.** *Am J Psychiatry* 2008, **165**(9):1155–1162.
68. Weissman MM, Pilowsky DJ, Wickramaratne PJ, Talati A, Wisniewski SR, Fava M, Hughes CW, Garber J, Malloy E, King CA, Cerda G, Sood AB, Alpert JE, Trivedi MH, Rush AJ, STAR*D-Child Team: **Remissions in maternal depression and child psychopathology: a STAR*D-child report.** *JAMA* 2006, **295**(12):1389–1398.

doi:10.1186/1471-244X-12-188

Cite this article as: Low *et al.*: The association between parental history of diagnosed mood/anxiety disorders and psychiatric symptoms and disorders in young adult offspring. *BMC Psychiatry* 2012 **12**:188.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

