

CORRECTION

Open Access



Correction to: Stability of personality traits over a five-year period in Swedish patients with schizophrenia spectrum disorder and nonpsychotic individuals: a study using the Swedish universities scales of personality

Tomas Fagerberg^{1*}, Erik Söderman¹, J. Petter Gustavsson², Ingrid Agartz^{1,3,4} and Erik G. Jönsson^{1,3}

Correction to: *BMC Psychiatry*

<https://doi.org/10.1186/s12888-018-1617-y>

After publication of the original article [1], it was brought to our attention that some of the numbers in Table 3 were incorrectly reported due to data entry errors. The corrected Table 3 can be found below - modified values are marked with bold. We apologize for the inconvenience this may have caused.

Author details

¹Human Brain Informatics (HUBIN), Department of Clinical Neuroscience, Centre for Psychiatric Research, Psychiatry Section, Karolinska Institutet and Hospital, Stockholm, Sweden. ²Division of Psychology, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden. ³NORMENT, KG Jebsen Centre for Psychosis Research, Institute of Clinical Medicine, Psychiatry Section, University of Oslo, Oslo, Norway. ⁴Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway.

Received: 19 March 2019 Accepted: 19 March 2019

Published online: 08 April 2019

Reference

1. Fagerberg, et al. Stability of personality traits over a five-year period in Swedish patients with schizophrenia spectrum disorder and nonpsychotic individuals: a study using the Swedish universities scales of personality. *BMC Psychiatry*. 2018;18:54. <https://doi.org/10.1186/s12888-018-1617-y>.

* Correspondence: Tomas.Fagerberg@ki.se

¹Human Brain Informatics (HUBIN), Department of Clinical Neuroscience, Centre for Psychiatric Research, Psychiatry Section, Karolinska Institutet and Hospital, Stockholm, Sweden

Full list of author information is available at the end of the article



Table 3 Swedish universities Scales of Personality (SSP) raw scores (mean [standard deviation]) and 95% confidence interval (CI) of the difference between baseline and follow-up in patients with psychotic illness and non-psychotic controls (Continued)

	Control	2.21 (0.48)	2.03 (0.49)	0.09–0.26	1.75 (0.50)	1.60 (0.47)	0.06–0.24
ALL	Men	2.26 (0.49)	2.04 (0.53)	0.12–0.31	1.97 (0.56)	1.86 (0.57)	0.02–0.21
	Women	2.24 (0.55)	2.11 (0.50)	–0.03 – 0.29	1.81 (0.59)	1.72 (0.58)	–0.03 – 0.21
PATIENT	Men	2.36 (0.56)	2.16 (0.53)	0.01–0.38	2.28 (0.58)	2.25 (0.53)	–0.14 – 0.21
	Women	2.32 (0.66)	2.04 (0.74)	–0.37 – 0.94	2.20 (0.62)	2.29 (0.55)	–0.24 – 0.07
CONTROL	Men	2.20 (0.45)	1.97 (0.51)	0.11–0.34	1.79 (0.47)	1.62 (0.47)	0.05–0.28
	Women	2.22 (0.53)	2.13 (0.43)	–0.05 – 0.22	1.70 (0.54)	1.56 (0.49)	–0.01 – 0.29
		Verbal Trait Aggression			Physical Trait Aggression		
		Baseline	Follow-up	CI	Baseline	Follow-up	CI
ALL	Patient	2.01 (0.64)	2.00 (0.55)	–0.19 – 0.20	1.71 (0.57)	1.84 (0.60)	–0.31 – 0.07
	Control	2.15 (0.47)	1.96 (0.46)	0.10–0.28	1.93 (0.57)	1.67 (0.54)	0.17–0.35
ALL	Men	2.12 (0.53)	2.02 (0.52)	–0.01 – 0.21	1.92 (0.60)	1.81 (0.59)	0.00–0.23
	Women	2.07 (0.55)	1.88 (0.43)	0.04–0.33	1.73 (0.53)	1.55 (0.46)	0.01–0.33
PATIENT	Men	2.03 (0.62)	2.07 (0.58)	–0.26 – 0.18	1.76 (0.59)	1.86 (0.66)	–0.33 – 0.13
	Women	1.95 (0.74)	1.79 (0.38)	–0.37 – 0.69	1.54 (0.47)	1.74 (0.39)	–0.57 – 0.16
CONTROL	Men	2.18 (0.46)	2.00 (0.48)	0.07–0.31	2.02 (0.58)	1.78 (0.55)	0.14–0.35
	Women	2.10 (0.50)	1.91 (0.44)	0.04–0.34	1.78 (0.54)	1.50 (0.47)	0.11–0.44